



Nutrition Factsheet

March 2018

connect to the world of dairy

DAIRY AND CHILDREN: A HEALTHY COMBINATION

The importance of milk and dairy products for children and adolescents

- › Dairy foods are **natural sources of valuable nutrients for children**: high quality protein, vitamins and minerals such as calcium, phosphorus and iodine.
- › Research shows that dairy has **beneficial effects on health** beyond its nutritional value.
- › **Dairy products have an important place in children's diet**. They contribute to a good physical and cognitive development, health maintenance and good eating habits for life.

DAIRY HELPS CHILDREN TO REACH ESSENTIAL NUTRIENTS DIETARY RECOMMENDATIONS

- Milk and dairy are **naturally nutrient-rich foods^a** because, while they provide relatively few calories, they offer a whole range of **essential nutrients** (e.g. high quality proteins) and make a significant contribution to the daily nutrient intakes for many **vitamins and minerals** such as vitamin B12 and B2, calcium, phosphorus and iodine(1-3).
- Dairy products are considered an important part of a healthy balanced diet (4) that contribute to a better nutrient status (5-7). Therefore, dietary advice all over Europe recommends a daily consumption of dairy products during all stages of life. The average recommendations are 3-4 servings of dairy per day (France, Belgium, Ireland, Spain) or 500-600 ml of dairy per day (Denmark, Finland, the Netherlands) (Figure 1).

EUROPEAN AVERAGE RECOMMENDATIONS OF DAIRY CONSUMPTION

Adults: 2-3 servings/day
Children: 3-4 servings/day

One serving of dairy:

Milk

Yoghurt

Cheese



150-200ml



125g



20-30g

Figure 1. Average recommendations of dairy intake.



- Dairy consumption during childhood plays a vital role. However, **children's overall milk intake is declining** (8,9) specially in developed countries through middle childhood and adolescence (10). According to the EsKiMo Study (11), in Germany, less than 50% of the children achieve the recommended daily amounts of dairy. Many children and adolescents do not reach recommendations for dairy, and this leads to insufficient intakes of nutrients such as **calcium, selenium, and iodine** (12,13). Recent observations show already public health effects of this decline (14-16).

^aMore in EDA nutrition factsheet "[Nutrient-rich dairy, an affordable source of nutrition](#)"



Nutrition Factsheet

connect to the world of dairy

- Including a wide range of milk and dairy foods as a core component of school meals helps children to achieve dietary recommendations and consume a healthy and nutrient-rich diet.

- Besides basic plain milk and dairy, studies show that having a wide variety of **flavoured milks and yoghurts can help to increase milk consumption** and boosting their vitamin, mineral and protein intake without any adverse impact on weight (17-20)^b. A variety of studies also show that a frequent consumption of yoghurt is related with a better diet overall (19,20,21). In some countries, **children that drink flavoured milks have a higher total milk intake** than those who drink regular milk, while intake of added sugars does not differ much between those groups (22). Children who drink both plain and flavoured milk have significantly higher intakes of vitamin A, calcium, phosphorus, magnesium, potassium than those who only drink plain milk (22). A US study showed that removing flavoured milks from schools leads to a decrease in overall milk consumption and negatively impacts children’s nutrients intake (23).



MILK AND DAIRY NUTRIENTS ARE IMPORTANT FOR CHILDREN’S HEALTH

- **Dairy nutrient-rich profile** contributes to a good development and functioning of the body and health maintenance from early ages (24) and across all life stages^c. During childhood and adolescence, **bones need high quality protein and calcium** to grow and develop healthily and also to maintain bone health later in life (25). **Both nutrients are naturally abundant in dairy products such as milk, cheese and yoghurts.** The health benefits of these products can be communicated to consumers through many authorised EU nutrition and health claims (26).
- Iodine, a nutrient that can be found in milk and cheese, plays an important role in children’s normal growth (16).

DID YOU KNOW...
 ...Dairy calcium is better absorbed than calcium from plant sources (27) and accounts for more than 50% of children’s total calcium intake in European countries (5).

- **Nutrients in dairy are also important for muscles.** High quality proteins in milk and dairy foods contains many essential amino acids and bioactive peptides which may have specific effects on growth (28) and contribute to muscle growth and maintenance (29).

- Many observational studies also suggest a positive association between dairy intake and **dental health**. Particularly cheese and yoghurt consumption leads to less dental caries in children (30,31). Concerning cardiovascular risks, scientific data shows that **higher dairy**



^bMore in EDA nutrition factsheet “[Sugar in dairy products](#)”

^cMore in EDA nutrition factsheet “[Health benefits and nutritional value of dairy](#)”



Nutrition Factsheet

connect to the world of dairy

intake in early life is associated with lower blood pressure in middle childhood (32) or early adolescence (33). There is also a positive association between yoghurt consumption, including flavoured ones, and diet quality and metabolic profile in children (34).

- Recent data shows that more and more children in Europe drink low-fat milk (10,35,36). This can alleviate the concern that energy provided by dairy may contribute to childhood obesity. Dairy products (excluding butter) account for about **10% of total fat intake** in German children aged 1-13 years (35). In the Netherlands, dairy products account for **25% of total energy**, while milk alone contributed **around 6% of total energy** to the diet of young children (6). Moreover, a recent review found **no association between dairy intake and measures of adiposity** (BMI, body fat) (37). Another study among adolescents in Portugal also found that those who consumed 2 cups or more of milk per day were significantly less likely to have abdominal obesity than those who consumed less milk (38).

DID YOU KNOW...

...Milk and dairy foods are a very affordable source of essential nutrients. Milk and yoghurt provide excellent nutrition value in relation to cost (1).



DRINKING MILK CONTRIBUTES TO HEALTHY EATING HABITS FOR A HEALTHY LIFE

- Milk and dairy are recommended to be part of breakfast at home and childcare facilities** (39). This helps children to develop the taste for milk from the childhood and to **adopt healthy eating habits** later in life, like **reaching dairy intake recommendations and having breakfast every day**. 10-30% of children and adolescents in Europe and in the USA skip breakfast, which may have unfavourable consequences for health and cognitive performance (39). Growing scientific evidence shows that there is a relation between skipping breakfast and nutritional inadequate diets as well as with obesity (40). Several studies found **higher milk consumption and calcium intakes in children and adolescents who eat breakfast** (41,42).



Dairy products are available in all kind of taste preferences



Nutrition Factsheet

connect to the world of dairy

- **Children’s overall milk intake is decreasing (8,9,35) and it is often replaced by potentially nutrient-poor, energy-dense foods and beverages (43,44) that contain “empty calories”.** Nutrient-rich foods such as dairy are important for a balanced and healthy diet (2), and replacing them with sweetened beverages leads to an increase in energy intake and to a decrease in calcium intake (45). Studies show a rise in childhood obesity coinciding with a decline in dairy consumption and an increase in sweetened, nutrient-poor, beverages consumption (46,47).





Nutrition Factsheet

connect to the world of dairy

REFERENCES

1. Drewnowski A, Fulgoni V. Nutrient profiling of foods: creating a nutrient-rich food index. *Nutr Rev* [Internet]. 2008 Feb;66(1):23–39. Available from: <https://academic.oup.com/nutritionreviews/article-lookup/doi/10.1111/j.1753-4887.2007.00003.x>
2. Streppel MT, de Groot LCPGM, Feskens EJM. Nutrient-rich foods in relation to various measures of anthropometry. *Fam Pract* [Internet]. 2012 Apr;29(suppl 1):i36–43. Available from: <https://academic.oup.com/fampra/article-lookup/doi/10.1093/fampra/cmr093>
3. Food and Agriculture Organization of the United Nations (FAO). Milk and dairy products in human nutrition [Internet]. Muehlhoff E, Bennett A, McMahon D, editors. Rome; 2013. 404 p. Available from: <http://www.fao.org/docrep/018/i3396e/i3396e.pdf>
4. Food and Agriculture Organization of the United Nations (FAO). Fats and fatty acids in human nutrition. Report of an expert consultation [Internet]. Geneva; 2008. Available from: <http://foris.fao.org/preview/2553-0ece4cb94ac52f9a25af77ca5cfba7a8c.pdf>
5. Coudray B. The Contribution of Dairy Products to Micronutrient Intakes in France. *J Am Coll Nutr* [Internet]. 2011 Oct;30(sup5):410S–414S. Available from: <http://www.tandfonline.com/doi/abs/10.1080/07315724.2011.10719984>
6. Vissers PA., Streppel MT, Feskens EJ., de Groot LCPG. The Contribution of Dairy Products to Micronutrient Intake in The Netherlands. *J Am Coll Nutr* [Internet]. 2011 Oct;30(sup5):415S–421S. Available from: <http://www.tandfonline.com/doi/abs/10.1080/07315724.2011.10719985>
7. Van Staveren WA, de Groot LCPGM. Evidence-based dietary guidance and the role of dairy products for appropriate nutrition in the elderly. *J Am Coll Nutr* [Internet]. 2011 Oct;30(5 Suppl 1):429S–37S. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22081689>
8. Lasater G, Piernas C, Popkin BM. Beverage patterns and trends among school-aged children in the US, 1989-2008. 2011; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3196913/pdf/1475-2891-10-103.pdf>
9. Lioret S, Dubuisson C, Dufour A, Touvier M, Calamassi-Tran G, Maire B, et al. Trends in food intake in French children from 1999 to 2007: results from the INCA (étude Individuelle Nationale des Consommations Alimentaires) dietary surveys. 2017; Available from: https://www.cambridge.org/core/services/aop-cambridge-core/content/view/21EBD680B892CAD3867ADD833B8F7A05/S0007114509992078a.pdf/trends_in_food_intake_in_french_children_from_1999_to_2007_results_from_the_inca_etude_individuelle_nationale_des_consommations
10. Bates B, Lennox A, Prentice A, Bates C. National Diet and Nutrition Survey: Headline results from Years 1,2 and 3 (combined) of the Rolling Programme (2008/2009-2010/11) [Internet]. London; 2012. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/207708/NDNS-Y3-report_All-TEXT-docs-combined.pdf
11. Mensink G, Heseke H, Richter A, Stahl A, Vohmann C. Ernährungsstudie als KiGGS-Modul (EsKiMo) [Internet]. 2007. Available from: https://www.bmel.de/SharedDocs/Downloads/Ernaehrung/EsKiMoStudie.pdf?__blob=publicationFile
12. Mouratidou T, Vicente-Rodriguez G, Gracia-Marco L, Huybrechts I, Sioen I, Widhalm K, et al. Associations of dietary calcium, vitamin D, milk intakes, and 25-hydroxyvitamin D with bone mass in Spanish adolescents: the HELENA study. *J Clin Densitom* [Internet]. 2013 Jan 1;16(1):110–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22901550>
13. Roman Viñas B, Ribas Barba L, Ngo J, Gurinovic M, Novakovic R, Cavelaars A, et al. Projected prevalence of inadequate nutrient intakes in Europe. *Ann Nutr Metab* [Internet]. 2011;59(2–4):84–95. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22142665>
14. Black RE, Williams SM, Jones IE, Goulding A. Children who avoid drinking cow milk have low dietary calcium intakes and poor bone health. *Journal of Clinical Nutrition* [Internet]. 2002;76(3):675–80. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12198017>
15. Rubio-Lopez N, Llopis-Gonzalez A, Morales-Suarez-Varela M. Calcium intake and nutritional adequacy in Spanish Children: The ANIVA Study. *Nutrients* [Internet]. 2017 Feb;9(2):170. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5331601/>
16. Kapil U. Health Consequences of Iodine Deficiency. *Sultan Kaboos Uni Med J* [Internet]. 2007 Dec;7(3):267–72. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5331601/>
17. Nicklas TA, O’Neil CE, Fulgoni VL. The Nutritional Role of Flavored and White Milk in the Diets of Children. *J Sch Health* [Internet]. 2013 Oct;83(10):728–33. Available from: <http://doi.wiley.com/10.1111/josh.12087>
18. Fayet F, Ridges LA, Wright JK, Petocz P. Australian children who drink milk (plain or flavored) have higher milk and micronutrient intakes but similar body mass index to those who do not drink milk. *Nutr Res* [Internet]. 2013 Feb;33(2):95–102. Available from:



Nutrition Factsheet

connect to the world of dairy

- <http://www.sciencedirect.com/science/article/pii/S0271531712002874?via%3Dihub>
19. Zhu Y, Wang H, Hollis J, Jaques P. The associations between yogurt consumption, diet quality, and metabolic profiles in children in the USA. *European Journal of Nutrition* [Internet]. 2015 Jun; 54(4): 543-50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25034879>
 20. Hobbs D, Givens D, Lovegrove J. Yogurt consumption is associated with higher nutrient intake, diet quality and favourable metabolic profile in children: a cross-sectional analysis using data from years 1-4 of the National diet and Nutrition Survey, UK. *European Journal of Nutrition* [Internet]. 2017 May. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29330662>
 21. Moreno L, Bel-Serrat S, Santaliestra-Pasias A, Bueno G. Dairy products, yogurt consumption, and cardiometabolic risk in children and adolescents. *Nutrition Reviews* [Internet]. 2015 Aug; 73 Suppl 1:8-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26175484>
 - 22.. Murphy MM, Douglass JS, Johnson RK, Spence LA. Drinking Flavored or Plain Milk Is Positively Associated with Nutrient Intake and Is Not Associated with Adverse Effects on Weight Status in US Children and Adolescents. *J Am Diet Assoc* [Internet]. 2008 Apr;108(4):631-9. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0002822308000059>
 - 23.. Quann EE, Adams D. Impact on Milk Consumption and Nutrient Intakes From Eliminating Flavored Milk in Elementary Schools. *Nutr Today* [Internet]. 2013;48(3):127-34. Available from: <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00017285-201305000-00007>
 24. Koletzko B, Armbruster M, Bauer C, Bos K, Cierpka M, Cremer M, et al. Ernährung und Bewegung im Kleinkindalter. Handlungsempfehlungen des Netzwerks "Gesund ins Leben – Netzwerk Junge Familie", ein Projekt von IN FORM. *Monatsschrift Kinderheilkd* [Internet]. 2013;16. Available from: http://shop.aid.de/_assets/downloads_free/3418_2013_he_kleinkind_x000.pdf
 25. Commission Regulation (EC) No 983/2009 of 21 October 2009 on the authorisation and refusal of authorisation of certain health claims made on food and referring to the reduction of disease risk and to children's development and health. [Internet]. (OJ L 277, 22.10.2009, p. 3-12). Available from: <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32009R0983&from=EN>
 26. EU Register of nutrition and health claims made on foods [Internet]. Available from: http://ec.europa.eu/food/safety/labelling_nutrition/claims/register/public/?event=register.home
 27. Miller GD, Jarvis JK, McBean LD. The importance of meeting calcium needs with foods. *J Am Coll Nutr* [Internet]. 2001 Apr;20(2 Suppl):168S-185S. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11349940>
 28. Hoppe C, Mølgaard C, Michaelsen KF. Cow's Milk and Linear Growth in Industrialized and Developing Countries. *Annu Rev Nutr* [Internet]. 2006 Aug 18;26(1):131-73. Available from: <http://www.annualreviews.org/doi/10.1146/annurev.nutr.26.010506.103757>
 29. Commission Regulation (EU) No 432/2012 of 16 May 2012 establishing a list of permitted health claims made on foods, other than those referring to the reduction of disease risk and to children's development and health [Internet]. (OJ L 136, 25.5.2012, p. 1-40). Available from: <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=OJ:L:2012:136:FULL&from=EN>
 30. Tanaka K, Miyake Y, Sasaki S. Intake of dairy products and the prevalence of dental caries in young children. *J Dent* [Internet]. 2010 Jul 1;38(7):579-83. Available from: <http://www.sciencedirect.com/science/article/pii/S0300571210000898?via%3Dihub>
 31. Głabska D, Sińska B, Remiszewski A. Analysis of the dependence between milk and dairy products consumption, and dental caries observed in group of children and teenagers. *Rocz Panstw Zakł Hig* [Internet]. 2007;58(1):69-75. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17711093>
 32. Rangan AM, Flood VL, Denyer G, Ayer JG, Webb KL, Marks GB, et al. The effect of dairy consumption on blood pressure in mid-childhood: CAPS cohort study. *Eur J Clin Nutr* [Internet]. 2012 Jun 11;66(6):652-7. Available from: <http://www.nature.com/doifinder/10.1038/ejcn.2011.218>
 33. Yuan WL, Kakinami L, Gray-Donald K, Czernichow S, Lambert M, Paradis G. Influence of dairy product consumption on children's blood pressure: results from the QUALITY cohort. *J Acad Nutr Diet* [Internet]. 2013 Jul 1;113(7):936-41. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23684662>
 34. Zhu Y, Wang H, Hollis JH, Jacques PF. The associations between yogurt consumption, diet quality, and metabolic profiles in children in the USA. *Eur J Nutr* [Internet]. 2015 Jun;54(4):543-50. Available from: <http://link.springer.com/10.1007/s00394-014-0735-7>
 35. Alexy U, Kersting M. Time trends in the consumption of dairy foods in German children and adolescents. *Eur J Clin Nutr* [Internet]. 2003 Oct 24;57(10):1331-7. Available from: <http://www.nature.com/articles/1601696>
 36. Huybrechts I, Matthyss C, Vereecken C, Maes L, Temme EHM, Van Oyen H, et al. Food intakes by preschool children in Flanders compared with dietary guidelines. *Int J Environ Res Public Health*. 2008;5(4):243-57.



Nutrition Factsheet

connect to the world of dairy

37. Dror DK, Allen LH. Dairy product intake in children and adolescents in developed countries: trends, nutritional contribution, and a review of association with health outcomes. *Nutr Rev* [Internet]. 2014 Feb 1;72(2):68–81. Available from: <https://academic.oup.com/nutritionreviews/article-lookup/doi/10.1111/nure.12078>
38. Abreu S, Santos R, Moreira C, Santos PC, Vale S, Soares-Miranda L, et al. Relationship of milk intake and physical activity to abdominal obesity among adolescents. *Pediatr Obes* [Internet]. 2014 Feb 1;9(1):71–80. Available from: <http://doi.wiley.com/10.1111/j.2047-6310.2012.00130.x>
39. Rampersaud GC, Pereira MA, Girard BL, Adams J, Metz J. Breakfast Habits, Nutritional Status, Body Weight, and Academic Performance in Children and Adolescents. *J Am Diet Assoc* [Internet]. 2005 May 1;105(5):743–60. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0002822305001513>
40. Deshmukh-Taskar PR, Nicklas TA, O'Neil CE, Keast DR, Radcliffe JD, Cho S. The Relationship of Breakfast Skipping and Type of Breakfast Consumption with Nutrient Intake and Weight Status in Children and Adolescents: The National Health and Nutrition Examination Survey 1999-2006. *J Am Diet Assoc* [Internet]. 2010 Jun 1;110(6):869–78. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0002822310002439>
41. Bowman SA. Beverage Choices of Young Females. *J Am Diet Assoc* [Internet]. 2002 Sep 1;102(9):1234–9. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0002822302902737>
42. Ortega RM, Requejo AM, López-Sobaler AM, Andrés P, Quintas ME, Navia B, et al. The importance of breakfast in meeting daily recommended calcium intake in a group of schoolchildren. *J Am Coll Nutr* [Internet]. 1998 Feb;17(1):19–24. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9477385>
43. Keller KL, Kirzner J, Pietrobelli A, St-Onge M-P, Faith MS. Increased Sweetened Beverage Intake Is Associated with Reduced Milk and Calcium Intake in 3- to 7-Year-Old Children at Multi-Item Laboratory Lunches. *J Am Diet Assoc* [Internet]. 2009 Mar 1;109(3):497–501. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0002822308021937>
44. Striegel-Moore RH, Thompson D, Affenito SG, Franko DL, Obarzanek E, Barton BA, et al. Correlates of beverage intake in adolescent girls: the National Heart, Lung, and Blood Institute Growth and Health Study. *J Pediatr* [Internet]. 2006 Feb 1;148(2):183–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16492426>
45. Yen ST, Lin B. Beverage consumption among US children and adolescents: full-information and quasi maximum-likelihood estimation of a censored system. *Eur Rev Agric Econ* [Internet]. 2002 Mar 1;29(1):85–103. Available from: <https://academic.oup.com/erae/article-lookup/doi/10.1093/erae/29.1.85>
46. Bray GA, Nielsen SJ, Popkin BM. Consumption of high-fructose corn syrup in beverages may play a role in the epidemic of obesity. *Am J Clin Nutr* [Internet]. 2004 Apr 1;79(4):537–43. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15051594>
47. Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr* [Internet]. 2006 Aug 1;84(2):274–88. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16895873>