



Milk Fat

Milk fat is complex and nutritionally unique

Dairy fat is the most complex fat source of human diet – it contains nearly 400 different fatty acids, many of them unique to dairy.



Nutrient richness of dairy includes its fatty acids as well as other nutrients

Dairy contains not only different fatty acids but is also a major source of high-quality proteins (whey and casein), many vitamins, minerals and much more!

Low and full-fat dairy is essential in a healthy and balanced diet

Dairy products, including regular-fat milk, cheese and yoghurt are part of a healthy and balanced diet as indicated in national dietary guidelines in all European countries.




Health effects of fatty acids depend on the food source

There is considerable evidence that the health effects of saturated fat vary depending on the specific fatty acid and on the specific food source.

I. Fat content of dairy products

Dairy is a diverse group of products with a broad variation in nutritional composition. Habitual intake patterns across European countries can also greatly vary. Dairy products with different fat levels have been available for a long time in all countries.






Milk and Yoghurt

Raw milk's fat content varies around 4%. Milk sold in stores is usually available as skimmed (below 0.5%), semi-skimmed (1.5-1.8%) and whole (3.5-4.0%). However, other fat levels are also available. Yoghurt and yoghurt drinks can be made from all types of milk and are therefore also available in different fat levels.

Cheese

Cheeses are available in many different forms - e.g. hard and soft cheeses, as spreads or slices - and with very different fat levels. In the EU, this level can amount up to around 45% depending on the type of cheese. [See average fat content of different types of cheeses](#) ▶▶▶



Cream

Cream is available in different fat levels (from 10% to around 35%). It is usually used for cooking and eaten together with other foods.

Butter

Butter is used as spread or cooking ingredient and consumed together with other foods such as bread and vegetables. The EU legislation recognises the fat content of milk and butter as a quality criterium; the current CMO Regulation ([n° 1308/2013](#)) defines "butter" as containing "not less than 80% but less than 90%" of milk fat".



II. Composition of milk fat

The main characteristic of milk fat is the variety of fatty acids it contains: more than 400 different fatty acids! [26]

Dairy products are defined by the raw material milk and their final nutrient content depends on the natural composition of the milk. The composition of milk fat varies slightly according to the breed of the cow, the stage of lactation, the season, the geographical location and the feed composition.


Milk fat fatty acids are divided into four main categories:

- Saturated Fatty Acids (SFA)
- Mono-Unsaturated Fatty Acids (MUFA)
- Poly-Unsaturated Fatty Acids (PUFA)
- Conjugated Linoleic (fatty) Acids (CLFA)

Milk fat contains on average 65-70% of saturated fatty acids (SFA) and 30-35% of mono (MUFA) and poly-unsaturated fatty acids (PUFA). [17]

Saturated Fatty Acids (SFA)

Typical fatty acid profile of cow's milk



There is considerable evidence that SFA health effects vary depending on the specific fatty acid and on the specific food source.



Food-based meta-analyses constantly find no association between dairy foods and increased risk of cardiovascular disease (CVD) [27, 35, 36, 35].



A recent large-scale study reported dairy consumption to be associated with lower risk of mortality and of major CVD events [8].

Other studies even found that whole-fat fermented dairy, e.g. cheese and yoghurts, may actually reduce CVD and diabetes risk [8, 19, 1, 9, 32, 33].

These results might be explained by the specificity of the interactions between dairy nutrients, i.e. different fatty acids, proteins (whey and casein), minerals (calcium, magnesium, phosphate), sodium, and phospholipid components of milk fat globule membrane (MFGM). This phenomenon is best known as the **dairy matrix**, meaning that the interactions between the structure of a food and the nutrient content of a product influences the whole digestion and absorption process. Indeed, nutrients are not eaten individually but as part of a whole food [38].



“Cheese paradox” explained by dairy matrix

Cheese composition has been shown to be more similar to yoghurt and milk than to butter, due to protein, mineral and MFGM contents even if it has higher contents of fat and salt. Furthermore, studies show that the interactions between nutrients present in cheese work in concert to reduce risks of CVD [39] showing that foods are not the sum of their nutrients but are part of a matrix [30].

Unsaturated Fatty Acids

Amongst unsaturated fatty acids in milk fat, there is around 25-30% MUFA among which oleic acid is the most important, and 2-6% PUFA, including in healthy proportions both omega 3 (that helps improving heart health, reducing weight in obese and overweight subjects, decreasing liver fat, and are anti-inflammatory) and omega 6 (that are precursor to other substances in human body, and are pro-inflammatory) fatty acids.

Trans Fatty Acids

Trans fatty acids (TFAs) are also unsaturated fatty acids. They can be industrially produced (e.g. frying oils, margarines, spreads) or occurring naturally in small quantities in meat and dairy.

Natural TFAs (also called ruminant TFAs) are naturally produced in the rumen (a special stomach that ferments plant-based foods prior to the digestion) of the cow due to a bacterial fermentation. Levels of natural TFAs found in dairy are very low, e.g. 0.09 g/100g in full fat milk, 0.03 g/100g in low fat yoghurt and 0.87 g/100g in cheddar cheese [14].

Since 2021, EU Regulation 2019/649 has limited the maximum allowed content of industrial TFAs in food at 2 g/100g of fat. **This Regulation excludes trans-fat naturally occurring in fat of animal origin, including milk fat**

The literature shows that consumption of ruminant TFAs from natural sources such as dairy, meat and butter is below 2 g/day in the average European diet [14, 6]. Contribution of ruminant TFAs to the overall energy intake in our usual diets is therefore not of any health relevance (further reading in our [EDA Q&A on TFAs](#)), since it is minimal and below the maximum level for total TFA intake of 1% of energy set by WHO [31]. There is a scientific consensus stating that the intake of TFAs via usually consumed amounts of dairy does not have negative effects on health [28].

Conjugated Linoleic Acid

Conjugated Linoleic Acid (CLA) is one of the ruminant TFAs. Science indicates that consumption of natural TFAs found in milk and dairy, in particular CLA and its precursor vaccenic acid (VA), may be linked to numerous beneficial health effects such as improved blood lipid profiles and decreased cholesterol absorption [13, 20]. Trans fatty acids in dairy, but especially CLA, have been associated with a reduction of body fat mass in overweight and obese subjects [21, 22]. Another ruminant-specific TFA, trans-palmitoleic acid, has been found to substantially reduce the risk of type 2 diabetes [10]; this acid is not produced by the human body [26], and can only be introduced to the diet with dairy products. Anti-cancer properties of ruminant CLA have also been shown in in vitro studies [12].

III. Health effects of milk fat

Milk fat is complex and contains several positive components or is consumed together with several nutrients that are beneficial to the health (e.g. proteins, essential fatty acids, CLA).

Reduced risk of cardiovascular disease (CVD)

The latest scientific evidence indicates that dairy foods have either beneficial or neutral effects on the risk of cardiometabolic diseases [29]. Studies also show that the consumption of regular fat dairy foods is not associated with an increased risk of cardiovascular disease [16].



Fermented dairy products, such as cheese and yoghurt, are associated with a lower risk of cardiovascular diseases [30], and high-fat dairy, non-fermented milk, butter or cream are not associated with stroke risk [29]. The recommendation to limit SFA intake -thought to increase CVD risk- has persisted despite increasing level of evidence to the contrary. Most recent meta-analyses of randomized trials and observational studies found no beneficial effects of reducing SFA intake on cardiovascular disease (CVD) and total mortality, and instead found protective effects against stroke (34, 35, 16).

The interaction between the numerous nutrients within a food enhance health properties of dairy products, thanks to the dairy matrix effect. As mentioned earlier, this might explain for example why cheese shows such healthy effects, despite its content of SFAs and salt [6].



Reduced risk of overweight and type 2 diabetes

Several pieces of research have demonstrated an inverse association between milk fat and body fat and the risk of obesity [4], and a non-linear but strong favourable association between yoghurt intake and lower risk of type 2 diabetes [3, 15].

Prevention of hypertension

High-quality studies also highlight the positive impact of dairy with respect to a lower hypertension risk and a reduced blood pressure, the latest constituting a major risk factor for development of stroke, coronary heart disease, heart failure, and end-stage renal disease [11], especially through the cis-MUFA contained in dairy foods [24]. This is why milk is part of the Dietary Approach to Stop Hypertension (DASH) diet, a US guideline, composed to specifically prevent/treat hypertension.



Dairy is much more than milk fat



In Western diets, dairy products are **the primary source of calcium** providing between 40-70% of the daily needs. They also provide **many other minerals** and contribute to the intake of **important vitamins** such as fat-soluble vitamins A and D, vitamins B2, B6, B12, and provide **high-quality protein**.

All dietary guidelines in every EU country recommend dairy as part of a healthy diet [40]. They are rich in many nutrients and are key contributors of several minerals, vitamins and high-quality protein. Beyond pure nutritional value, several other health benefits have been demonstrated or suggested for all dairy foods [18].

Sources

- 1- Astrup A., *Yogurt and dairy product consumption to prevent cardiometabolic diseases: epidemiologic and experimental studies*, The American Journal of Clinical Nutrition, Volume 99, Issue 5, May 2014, Pages 1235S–1242S [2014].
<https://doi.org/10.3945/ajcn.113.073015>
- 2- Astrup Arne, Bertram Hanne CS, Bonjour Jean-Philippe, de Groot Lisette CP, de Oliveira Otto Marcia C, Feeney Emma L et al. *WHO draft guidelines on dietary saturated and trans fatty acids: time for a new approach?* BMJ 2019; 366 :l4137 [2013] <https://doi.org/10.1136/bmj.l4137>
- 3- Aune D, Norat T, Romundstad P, Vatten LJ. *Dairy products and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies*. Am J Clin Nutr 2013; 98: 1066–1083 [2013]
<https://doi.org/10.3945/ajcn.113.059030>
- 4- Astrup A. et al, Regular-Fat Dairy and Human Health: A Synopsis of Symposia Presented in Europe and North America (2014–2015), *Nutrients* 2016, 8(8), 463 [2016]. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4997376/>
- 5- Benbrook CM, Butler G, Latif MA, Leifert C, Davis DR. *Organic production enhances milk nutritional quality by shifting Fatty Acid composition: a United States-wide, 18-month study*. PLoS One. 2013 Dec 9;8(12):e82429. DOI: [10.1371/journal.pone.0082429](https://doi.org/10.1371/journal.pone.0082429)
- 6- Craig-Schmidt MC. World-wide consumption of trans fatty acids. *Atheroscler Suppl*. 2006 May;7(2):1–4
doi:[10.1016/j.atherosclerosissup.2006.04.001](https://doi.org/10.1016/j.atherosclerosissup.2006.04.001)
- 7- De Goede J, Geleijnse JM, Ding EL, Soedamah-Muthu SS. *Effect of cheese consumption on blood lipids: a systematic review and meta-analysis of randomized controlled trials*. Nutr Rev. 2015;73(5):259–75. [2015] DOI:
<https://doi.org/10.1093/nutrit/nuu060>
- 8- Dehghan M, Mente A, Rangarajan S et al., *Association of dairy intake with cardiovascular disease and mortality in 21 countries 5from five continents (PURE): a prospective cohort study*. The Lancet. 2018 Sept 11. PMID: 30217460 DOI:
[10.1016/S0140-6736\(18\)31812-9](https://doi.org/10.1016/S0140-6736(18)31812-9) [2018]
- 9 - de Oliveira Otto MC, Mozaffarian D, Kromhout D, et al., *Dietary intake of saturated fat by food source and incident cardiovascular disease: the Multi-Ethnic Study of Atherosclerosis*. Am J Clin Nutr. 2012;96(2):397–404.
doi:10.3945/ajcn.112.037770 [2012]. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3396447/>
- 10- de Souza Russell J, Mente Andrew, Maroleanu Adriana, et al. *Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies* BMJ 2015; 351 :h3978 [doi: 10.1136/bmj.h3978](https://doi.org/10.1136/bmj.h3978)
- 11- Drouin-Chartier JP et al., *Systematic Review of the Association between Dairy Product Consumption and Risk of cardiovascular-Related Clinical Outcomes*. Adv Nutr. 2016 Nov 15;7(6):1026–1040. [2016] doi: [10.3945/an.115.011403](https://doi.org/10.3945/an.115.011403).
- 12- Erickson K.L., Hubbard N.E., Borowsky A.D., Damonte P. (2010) *Conjugated Linoleic Acid and Cancer*. In: Milner J.A., Romagnolo D.F. (eds) *Bioactive Compounds and Cancer. Nutrition and Health*. Humana Press, Totowa, NJ. [2010] DOI https://doi.org/10.1007/978-1-60761-627-6_12
- 13- Gayet-Boyer C, Tenenhaus-Aziza F, Prunet C, et al. *Is there a linear relationship between the dose of ruminant trans-fatty acids and cardiovascular risk markers in healthy subjects: results from a systematic review and meta-regression of randomised clinical trials?* British Journal of Nutrition, Dec 2014, pp 1914–1922 DOI: [10.1017/S0007114514002578](https://doi.org/10.1017/S0007114514002578)
- 14- Gebauer S, Chardigny J-M., Uhre M, et al., *Effects of Ruminant trans Fatty Acids on Cardiovascular Disease and Cancer: A Comprehensive Review of Epidemiological, Clinical, and Mechanistic Studies*. Adv Nutr July 2011 Adv Nutr vol. 2: 332–354, 2011, [doi: 10.3945/an.111.000521](https://doi.org/10.3945/an.111.000521)

- 15- Gijsbers L, Ding EL, Malik VS, de Goede J, Geleijnse JM, Soedamah-Muthu SS. *Consumption of dairy foods and diabetes incidence: a dose-response meta-analysis of observational studies*. Am J Clin Nutr 2016;103:1111–24. [2016] DOI: <https://doi.org/10.3945/ajcn.115.123216>
- 16- Givens D. I., *Saturated Fats, dairy foods and health: A curious paradox?*, Nutr Bull, 42: 274-282. doi:10.1111/nbu.12283 [2017] DOI: <https://doi.org/10.1111/nbu.12283>
- 17- Givens D. I. and Parodi P., *Proceedings of the Symposium 'Scientific Update on Dairy Fats and Cardiovascular Diseases'*, Supplement to The Journal of the American College of Nutrition Volume 27 Number 6 December 2008 Pages 720S-754S [2008]
- 18- Godos J., Tieri M., Ghelfi F. *et al.* (2019) Dairy foods and health: an umbrella review of observational studies, International Journal of Food Sciences and Nutrition, DOI: [10.1080/09637486.2019.1625035](https://doi.org/10.1080/09637486.2019.1625035)
- 19- Ibsen DB, Laursen ASD, Lauritzen L, Tjønneland A, Overvad K, Jakobsen MU. *Substitutions between dairy product subgroups and risk of type 2 diabetes: the Danish Diet, Cancer and Health cohort*. British Journal of Nutrition, 2017;118:989-997. doi: 10.1017/S0007114517002896. 11. DOI : [10.1017/S0007114519000564](https://doi.org/10.1017/S0007114519000564)
- 20- Kuhnt K, Degen C, Jahreis G. Evaluation of the Impact of Ruminant trans Fatty Acids on Human Health: Important Aspects to Consider, 2015 DOI: [10.1080/10408398.2013.808605](https://doi.org/10.1080/10408398.2013.808605)
- 21- Labonté ME *et al.*, *Impact of dairy products on biomarkers of inflammation: a systematic review of randomized controlled nutritional intervention studies in overweight and obese adults*, The American Journal of Clinical Nutrition, Volume 97, Issue 4, April 2013, Pages 706–717 [2013] DOI : <https://doi.org/10.3945/ajcn.112.052217>
- 22- Lalithadevi B., Muthiah Ns., Satya Narayana Murty K. Antioxidant activity of conjugated linoleic acid. November 2018 Asian Journal of Pharmaceutical and Clinical Research 11(11):169 DOI: [10.22159/ajpcr.2018.v11i11.27700](https://doi.org/10.22159/ajpcr.2018.v11i11.27700)
- 23- Legrand P. *Intérêt nutritionnel des principaux acides gras des lipides laitiers*. Sciences des Aliments 2008. 28:34-43 [2008] DOI : [10.3166/sda.28.34-43](https://doi.org/10.3166/sda.28.34-43)
- 24- Liu X, Kris-Etherton PM, West SG *et al.* *Effects of Canola and high-oleic-acid Canola oils on abdominal fat mass in individuals with central obesity*. Obesity 24 : 2261 – 8. [2016] DOI : [10.1002/oby.21584](https://doi.org/10.1002/oby.21584)
- 25- Månsson H.L., *Fatty acids in bovine milk fat*. Food & nutrition research vol. 52 (): 10.3402/fnr.v52i0.1821. doi: [10.3402/fnr.v52i0.1821](https://doi.org/10.3402/fnr.v52i0.1821) [2008]
- 26- Mozaffarian D. *et al.* *Trans-Palmitoleic Acid, Metabolic Risk Factors, and New-Onset Diabetes in US Adults*, PMC 2011 Dec 21. DOI: [10.1059/0003-4819-153-12-201012210-00005](https://doi.org/10.1059/0003-4819-153-12-201012210-00005)
- 27- Pimpin L, Wu JHY, Haskelberg H, Del Gobbo L, Mozaffarian D. *Is butter back? A systematic review and meta-analysis of butter consumption and risk of cardiovascular disease, diabetes, and total mortality*. PLoSONE2016;11:e0158118.doi:10.1371/journal.pone.0158118. DOI : [10.1371/journal.pone.0158118](https://doi.org/10.1371/journal.pone.0158118)
- 28- Rioux V, Legrand P. *Saturated fatty acids: simple molecular structures with complex cellular functions*. Curr Opin Clin Nutr Metab Care 2007. 10(6):752-8. Willett W, Mozaffarian D. Ruminant or industrial sources of trans fatty acids: public health issue or food label skirmish? Am J Clin Nutr 2008. 87:515-516 [2007] DOI : [10.1097/MCO.0b013e3282f01a75](https://doi.org/10.1097/MCO.0b013e3282f01a75)
- 29- Soedamah-Muthu SS. & de Goede J., *Dairy Consumption and Cardiometabolic Diseases: Systematic Review and Updated Meta-Analyses of Prospective Cohort Studies*, J. Curr Nutr Rep (2018) 7: 171. [2018] DOI : <https://doi.org/10.1007/s13668-018-0253-y>
- 30- Thorning TK *et al.*, *Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps*. Am J Clin Nutr. 2017 May;105(5):1033-1045. DOI : [10.3945/ajcn.116.151548](https://doi.org/10.3945/ajcn.116.151548)

- 31- WHO. Diet, [Nutrition and the Prevention of Chronic Diseases. Report of a Joint WHO/FAO Expert Consultation](#). WHO Technical Report Series No. 916. World Health Organization: Geneva, 2003.
- 32- Hess JM, Cifelli CJ, Fulgoni VL 3rd. Modeling the Impact of Fat Flexibility With Dairy Food Servings in the 2015-2020 Dietary Guidelines for Americans Healthy U.S.-Style Eating Pattern. *Front Nutr*. 2020 Oct 22;7:595880. doi: 10.3389/fnut.2020.595880. PMID: 33195378; PMCID: PMC7642593.
- 33- Hirahatake KM, Astrup A, Hill JO, Slavin JL, Allison DB, Maki KC. Potential Cardiometabolic Health Benefits of Full-Fat Dairy: The Evidence Base. *Adv Nutr*. 2020 May 1;11(3):533-547. doi: 10.1093/advances/nmz132. PMID: 31904812; PMCID: PMC7231591.
- 34- Kang Z, Yang Y, Xiao B. Dietary saturated fat intake and risk of stroke : Systematic review and dose-response meta-analysis of prospective cohort studies. *Systematic Reviews and Meta-Analyses*, 2020 Feb, 30(2):179-189. <https://doi.org/10.1016/j.numecd.2019.09.028>
- 35- Astrup A, Magkos F, Bier DM, Brenna JT, de Oliveira Otto MC, Hill JO, King JC, Mente A, Ordovas JM, Volek JS, Yusuf S, Krauss RM. Saturated Fats and Health: A Reassessment and Proposal for Food-Based Recommendations: JACC State-of-the-Art Review. *J Am Coll Cardiol*. 2020 Aug 18;76(7):844-857. doi: 10.1016/j.jacc.2020.05.077. Epub 2020 Jun 17. PMID: 32562735.
- 36- Trieu K, Bhat S, Dai Z, Leander K, Gigante B, Qian F, Korat AVA, Sun Q, Pan XF, Laguzzi F, Cederholm T, de Faire U, Hellénus ML, Wu JHY, Risérus U, Marklund M. Biomarkers of dairy fat intake, incident cardiovascular disease, and all-cause mortality: A cohort study, systematic review, and meta-analysis. *PLoS Med*. 2021 Sep 21;18(9):e1003763. doi: 10.1371/journal.pmed.1003763. PMID: 34547017; PMCID: PMC8454979.
- 37- Astrup A, Teicholz N, Magkos F, Bier DM, Brenna JT, King JC, Mente A, Ordovas JM, Volek JS, Yusuf S, Krauss RM. Dietary Saturated Fats and Health: Are the U.S. Guidelines Evidence-Based? *Nutrients*. 2021 Sep 22;13(10):3305. doi: 10.3390/nu13103305. PMID: 34684304; PMCID: PMC8541481.
- 38 - Weaver, Connie M. "Dairy matrix: is the whole greater than the sum of the parts?." *Nutrition reviews* vol. 79,Suppl 2 (2021): 4-15. doi:10.1093/nutrit/nuab081
- 39 - Feeney, E.L., Lamichhane, P. and Sheehan, J.J. (2021), The cheese matrix: Understanding the impact of cheese structure on aspects of cardiovascular health – A food science and a human nutrition perspective. *Int J Dairy Technol*, 74: 656-670. <https://doi.org/10.1111/1471-0307.12755>
- 40 – European Commission, 2024, Health Promotion and Disease Prevention Knowledge Gateway – Food-Based Dietary Guidelines in Europe, Accessible here.